

# Characterization of Microsatellite Loci and Repeat Density in the Gooseneck Barnacle, *Pollicipes elegans*, Using Next Generation Sequencing

LOUIS V. PLOUGH AND PETER B. MARKO

From the Department of Biological Sciences, Clemson University, 132 Long Hall, Clemson 29634, South Carolina.

Address correspondence to Louis V. Plough, e-mail: [lpough@umces.edu](mailto:lpough@umces.edu).

Data deposited at Dryad: <http://dx.doi.org/10.5061/dryad.b7855>

*Pollicipes elegans* is a commercially important and biogeographically significant rocky-shore gooseneck barnacle found along the eastern Pacific coasts of Peru, El Salvador, and Mexico. Little is known about its reproductive biology, and no genetic resources exist despite its growing importance as a fisheries species in the region. Next generation sequencing methods can provide rapid and cost-effective development of molecular markers such as microsatellites, which can be applied to studies of paternity, parentage, and population structure in this understudied species. Here, we used Roche 454 pyrosequencing to develop microsatellite markers in *P. elegans* and made genomic comparisons of repeat density and repeat class frequency with other arthropods and more distantly related taxa. We identified 13 809 repeats of 1–6 bp, or a density of 9744 bp of repeat per megabase queried, which was intermediate in the range of taxonomic groups compared. Comparison of repeat class frequency distributions revealed that *P. elegans* was most similar to *Drosophila melanogaster* rather than the more closely related crustacean *Daphnia pulex*. We successfully isolated 15 polymorphic markers with an average of 9.4 alleles per locus and average observed and expected heterozygosities of 0.501 and 0.597, respectively. Four loci were found to be out of Hardy–Weinberg equilibrium, likely due to the presence of null alleles. A preliminary population genetic analysis revealed low but significant differentiation between a Peruvian ( $n = 47$ ) and Mexican ( $n = 48$ ) population ( $F_{ST} = 0.039$ ) and markedly reduced genetic diversity in Peru. These markers should facilitate future studies of paternity, parentage, and population structure in this species.

**Key words:** 454, barnacles, microsatellite, null alleles, paternity, population genetics

*Pollicipes elegans* is a gooseneck barnacle (order Pedunculata, class Pollicipea) that inhabits the rocky intertidal of the eastern

Pacific. *Pollicipes elegans* is found along a large latitudinal range spanning the equator, but with a disjunct distribution of populations in Northern Peru, El Salvador, and Northwest Mexico (Van Syoc 1994; Barnes 1996). The fleshy peduncle is an edible and highly prized seafood item from a subsistence fishery that has existed in parts of Peru for generations (Bernard 1988; Kameya and Zeballos 1988). A commercial fishery was developed in the 1970s to supply barnacles to the European market, after stocks of *Pollicipes pollicipes* in Portugal and Spain crashed due to overexploitation (Molares and Freire 2003). Overharvesting is now a concern for *P. elegans* in Peru, but it is unclear if there has been a significant reduction in biomass and genetic diversity due to fishing pressure. Few published studies of any kind exist for *P. elegans* (Barnes 1996), and only a single genetic study has been undertaken, which found relatively high divergence at the mitochondrial cytochrome oxidase 1 (CO1) locus between individuals from across the equator (Peru and Baja Mexico; Van Syoc 1994). Little is known about population structure among all populations and to what extent harvesting has affected genetic diversity in the region. Cryptic species may also exist for *P. elegans*, as the disjunct populations across the equator likely have limited gene flow between them. In the better-studied *P. pollicipes*, the finding of substantial genetic heterogeneity across populations ranging from North Africa to Northwest Europe (Quinteiro et al. 2007; Campo et al. 2010) led to the discovery that the isolated Cape Verdean population was a distinct species (Fernandes et al. 2010; Quinteiro et al. 2011).

In addition to the lack of population genetic information, there is a paucity of data on the basic reproductive biology and mating system of *P. elegans*, such as whether or not females store sperm, mate with multiple males (polyandry), or even spermcast (Bishop and Pemberton 2006). The question of paternity is equally open for all species of *Pollicipes*, and barnacles in general, because to date, only 1 study of paternity has been published for any barnacle species (Kelly et al. 2012).

The development of species-specific, highly polymorphic nuclear markers, such as microsatellites, would facilitate the analysis of multiple paternity (how many fathers contribute to a brood), studies of relatedness within barnacle aggregations (which can be high; Veliz et al. 2006), as well as standard and coalescent-based population genetic analysis of genetic divergence, divergence time, migration (current and historical), and diversity (e.g., effective population size) in *P. elegans*.

Microsatellites, highly variable genetic markers made up of short (1–6bp) tandem repeats, are the markers of choice for population and evolutionary genetic studies because of their codominant inheritance, high variability, high abundance in the genome, and relative ease of genotyping on a variety of platforms. Despite the increased popularity of single nucleotide polymorphisms as genetic markers, individual microsatellite markers consistently show greater power in analyses of population genetic structure, association studies, and sibling reconstructions (Ohashi and Tokunaga 2003; Rosenberg et al. 2003; Liu et al. 2005; Wang and Santure 2009); thus, they are still widely used across a range of taxa and experiments. Their isolation, detection, and development can, however, be relatively time consuming and expensive (e.g., Hamilton et al. 1999; Zane et al. 2002). With recent advances in next generation sequencing (NGS), 1000s to millions of short reads of random genomic sequence can be obtained quickly and at relatively low cost, and programs that screen for microsatellites in silico can be used to find repeat sequences and design suitable primers for polymerase chain reaction (PCR) amplification (e.g., Abdelkrim et al. 2009; Gardner et al. 2011). The large number of reads produced on NGS platforms also provide a snapshot of the genome, and with these data, one can investigate the genomic frequency and distribution of repeat types, providing important insights into the origin and evolution of microsatellites, and genome structure and organization more generally (Tóth et al. 2000; Ellergren 2004; Megléc et al. 2012a). No genome-wide repeat data have been generated for any barnacle species, so this analysis provides the first data of this sort for future genomic comparisons across Cirripedia.

Here, we use NGS on the Roche 454 FLX platform to develop and characterize microsatellite loci for use in future studies of paternity, parentage, and population structure in *P. elegans*. We also take advantage of the large amounts of sequence data produced with this shotgun sequencing approach and make genomic comparisons of the repeat density and diversity in *P. elegans* with other arthropods and more distantly related animal taxa, with the expectation that more closely related taxa will exhibit similar patterns of microsatellite density and repeat class frequency. Finally, we perform a preliminary analysis of genetic diversity and genetic structure in 2 populations to show the utility of these markers for population genetic analyses.

## Methods

### Samples and DNA Extractions

Barnacles from near the southern tip of the Baja California peninsula, Mexico (Gaspareno: 23°10'58.09"N, 110°8'26.51"W)

and Northwestern Peru (Islilla: 5°13'17.85"S, 81°11'18.99"W) were collected in 2009 and 2010, and peduncle tissue was stored in 95% ethanol. DNA was extracted using a modified hexadecyltrimethylammonium bromide (CTAB) protocol (Doyle and Dickson 1987), with 2 chloroform extraction steps and two 70% ethanol washes.

### 454 Sequencing and Quality Control

Shotgun sequencing of a genomic DNA library from a single individual from southern Baja, Mexico was performed in a portion of a single lane run on a 454 FLX Titanium sequencer (Roche 454 Life Sciences, Brandford, CT, USA), at the Genome Sequencing and Analysis Facility at the University of Texas at Austin. A total of 200 000 reads were requested, and library preparation proceeded following the manufacturer's protocol (Margulies et al. 2005). Briefly, DNA was fragmented through nebulization, and fragments in the 500–800-bp range were ligated to adaptors, captured on beads, and clonally amplified through emulsion PCR. After sequencing, raw sequences were filtered and trimmed for quality and length using the FASTQ tool kit on the Galaxy public server (<http://main.g2.bx.psu.edu>; Blankenberg et al. 2010). Reads were discarded if they were less than 100bp in length or if they had a mean quality score lower than 30. Low-quality bases from the 3' end of the read were trimmed with a sliding window, discarding bases until at least 2 individual scores were above a quality score of 35 (PHRED). After quality trimming, we performed a de novo assembly of reads using NEWBLER version 2.8 (Roche 454 Life Sciences) to eliminate duplicate reads, which can account for up to 25% of total reads from 454 FLX Titanium run (Niu et al. 2010; Balzer et al. 2013) and can bias upward the quantification of microsatellites in analyses of repeat density. After assembly using default parameters, contigs and unassembled reads were retained for repeat density analysis.

### Repeat Density Analysis and Microsatellite Discovery

To quantify the number and frequency of repeat classes and motifs throughout the sequences, we used the repeat scanning software PHOBOS version 3.3.12 (described in Mayer et al. 2010), which can search for perfect and imperfect microsatellites, giving a more accurate picture of repeat density across the genome compared with other repeat finding software. Using the PHOBOS results, we compared the genomic density of repeats and the frequency of each repeat class in *P. elegans* with genomic repeat data from a variety of animal species, including relatively closely related arthropods and the recently sequenced crustacean *Daphnia pulex*. Because repeat discovery rates can vary depending upon the software platform and the search parameters used, we used data from Mayer et al. (2010) and employed identical search parameters for *P. elegans* in PHOBOS version 3.3.12 to compare repeat data across taxa. Though our shotgun 454 reads represent only a fraction of the *P. elegans*' genome, inferences of genome-wide microsatellite frequency and density from low-coverage 454 reads have been shown to be highly correlated

with, and not significantly different from, estimates made from full, assembled genome data (Megléczy et al. 2012b).

To isolate microsatellites and design primers for population genetic analysis, we used the program QDD2.1 (beta) on the Linux platform (Megléczy et al. 2010). QDD2.1 searches for perfect microsatellites with repeat units of 1–6 bp and is able to filter for unique sequences and discard those associated with duplications or mobile elements. Default primer design settings were used except for the minimum flanking distance between a primer and a microsatellite (set to 30 bp) and the minimum PCR product size (set to 90 bp).

### Primer Testing

A total of 45 primer pairs from the QDD2.1 output were chosen for further development and analysis in *P. elegans*. Eight Mexican individuals were used initially to test for amplification success by visualizing amplification products on 1.0% low-melt agarose gels (Fisher Scientific, Waltham, MA, USA). Loci that amplified consistently were then genotyped in the 95 individuals sampled from Peru ( $n = 47$ ) and Mexico ( $n = 48$ ) to evaluate polymorphism and population genetic parameters. For genotyping, a 5' sequence tag that was complementary to a fluorescent probe was added either to the forward or to the reverse primer (FAM tag: GGAAACAGCTATGACCAT; VIC tag: CAGTCGGGCGTCATCA; NED tag: ACCAACCTAGGAAACACAG; and PET tag: GGCTAGGAAAGGTTAGTGGC; see Table 2 and Hauswaldt and Glenn 2003). In some cases, one of the 2 primers was directly labeled with a fluorophore if a suitable sequence tag could not be placed on the primer (see Table 2). PCR reactions were carried out in volumes of 12.5  $\mu$ L, with 0.5  $\mu$ L of each 5 mM primer (if direct labeled) or 0.05  $\mu$ L of the tagged primer, 0.45  $\mu$ L of fluorophore, and 0.5  $\mu$ L of reverse primer (if tag labeled: all primers at 5 mM), 2.5  $\mu$ L 5 $\times$  GoTaq buffer with 25 mM MgCl<sub>2</sub>, 0.3  $\mu$ L of 5 mM dNTPs, 0.125  $\mu$ L of 1% Triton-X (Fisher Scientific), and 0.1  $\mu$ L of GoTaq (Promega, Madison, WI, USA). Amplification was carried out separately for each locus on an MJ research Dyad thermocycler (now Bio-Rad, Hercules, CA, USA). Cycle conditions were as follows: an initial 2-min denaturing step at 94 °C, followed by 35 cycles of 30 s at 94 °C, 30 s at 53–57 °C (locus dependent; Table 2), and a 45-s extension step at 72 °C, all followed by a final extension step of 15 min at 72 °C. Successfully amplified loci were also tested for cross amplification in *P. pollicipes*, *P. polymerus*, and the recently described *P. caboverdensis/darwini* (Fernandes et al. 2010; Quinteiro et al. 2011). All sequencing was carried out at the DNA laboratory at Arizona State University. Genotypes were scored by eye using LIZ600 as an internal size standard on the Peak Scanner 1.0 software (Life Technologies, Carlsbad, CA, USA).

### Characterization of Novel Microsatellite Markers

We estimated overall number of alleles and observed and expected heterozygosities for the newly developed markers using FSTAT version 2.9.3.2 (Goudet 1995). Tests for Hardy–Weinberg equilibrium (HWE) were also carried out in FSTAT using 1500 permutations, and *P* values were

adjusted with a sequential Bonferroni correction. For those markers that differed significantly from HWE, estimates of null allele frequency were made using the FreeNA software (Chapuis and Estoup 2007). Pairwise tests for linkage disequilibrium (LD) were performed using 10 000 permutations in GENETIX (Belkhir et al. 2004).

### Analysis of Genetic Diversity and Population Structure in Peru and Mexico

We used the novel microsatellite markers to perform a preliminary population genetic analysis of the 2 populations sampled (Islilla, Peru and Gaspareno, Baja California, Mexico). Allelic richness was calculated using rarefaction in FSTAT version 2.9.3.2, and significant differences between the 2 populations were tested using the sign test (e.g., Jin et al. 2000). Gene diversity was also calculated in FSTAT, and significant differences between populations were assessed with a Wilcoxon signed-rank test (see Nei 1987, p. 183). To determine the degree of population differentiation, we estimated  $F_{ST}$  by calculating  $\Theta$  (Weir and Cockerham 1984), which accounts for small and unequal sample sizes, in FSTAT. Confidence intervals (95% CIs) for  $\Theta$  were calculated by bootstrapping across loci and used to assess overall significance.



## Results and Discussion

### 454 Sequencing Results

We obtained 191 746 reads with a mean length of 325 bp from a shared lane run on the Roche 454 FLX sequencing platform (Table 1). All reads have been deposited on the sequence read archive (accession SRP029161; <http://www.ncbi.nlm.nih.gov/sra>). Raw reads were then filtered and trimmed for length and quality, which resulted in the exclusion of 82 331 reads (47%), leaving 109 417 for downstream analysis. Cleaned reads were then assembled using NEWBLER version 2.8 into 973 contigs (maximum contig length of 8272 bp) and 99 324 unassembled reads, producing a set of 100 296 unique sequences for repeat analysis. These 100 296 sequences (36 805 630 nucleotides) accounted for approximately 2.83% of the genome of *P. elegans*, assuming a genome size of 1.3 gigabases, which was estimated from

**Table 1.** Raw read length statistics from Roche 454 sequencing

Median read length	358
Mean read length	325.07
Length	Number of sequences
100	16 267
200	17 536
300	26 544
400	72 831
500	56 240
600	2308
700	20
Total	191 746

**A****B**

**Figure 1.** (A) Genomic density of 1–6-bp repeat motifs for *Pollicipes elegans* and 12 other taxa versus genome size (log transformed). (B) Percentage distribution of repeat classes for each species. Following each taxon name on the x axis is the build, release, or version number of the genome used for the repeat analysis (for more information, see Table 1, Mayer et al. 2010).

the average sizes of 9 barnacle genomes from the animal genome database (Gregory 2012).

#### Genome-Wide Microsatellite Characterization

We found 13 809 repeats of 1–6bp comprising 358 619bp of total repeat sequence. We calculated that *P. elegans* has a repeat density of 9744bp of microsatellite repeats per

megabase (Mbase), which is slightly higher than that of the recently sequenced crustacean, *D. pulex* (7879 bp/Mbase) and similar to the fruit fly *Drosophila melanogaster* (10 875 bp/Mbase; Figure 1A). Repeat density appeared to scale somewhat with genome size, but this relationship was not significant (Pearson's correlation: 0.46,  $P = 0.11$ ). Comparing the frequency (bp/Mbase) of repeat classes in *P. elegans*, pentanucleotide repeats were the most common (29%), above that of

**Table 2.** Summary marker statistics and population genetic data

Marker data								Population genetic data					
Locus	GenBank number	Repeat	Allele number	Allele size <sup>a</sup>	$H_o^b$	$H_e$	Null <sup>c</sup>	$H_e$ Baja <sup>d</sup>	$H_e$ Peru	AR Baja <sup>e</sup>	AR Peru	$F_{ST}^f$	
1	Pole01	KF562712	AC	16	160–195	0.72	0.82		0.86	0.79	15.32	9.63	–0.001
2	Pole07	KF562713	AC	15	108–145 <sup>i</sup>	0.51	0.80	0.14	—	—	—	—	—
3	Pole08	KF562714	GAAT	8	141–177	0.41	0.49		0.56	0.41	6.48	6.34	0.007
4	Pole16	KF562715	AC	3	209–213 <sup>ii</sup>	0.25	0.28		0.34	0.22	3.00	2.00	0.001
5	Pole18	KF562716	TG	7	136–148 <sup>iii</sup>	0.50	0.51		0.52	0.49	5.83	5.82	0.006
6	Pole25	KF562717	GGC	5	203–215 <sup>i</sup>	0.53	0.57		0.64	0.53	7.31	3.73	0
7	Pole28	KF562718	GCAC	2	173–177 <sup>iii</sup>	0.17	0.28	0.04	—	—	—	—	—
8	Pole29	KF562719	CCCAGA	22	140–210	0.83	0.86		0.87	0.77	8.94	17.41	0.097
9	Pole30	KF562720	TG	6	149–159 <sup>ii</sup>	0.27	0.67	0.19	—	—	—	—	—
10	Pole34	KF562721	TGA	4	109–124 <sup>iv</sup>	0.63	0.67		0.66	0.41	3.97	3.00	0.09
11	Pole41	KF562722	AG	19	108–180 <sup>ii</sup>	0.59	0.80	0.10	—	—	—	—	—
12	Pole43	KF562723	AC	6	117–129 <sup>iv</sup>	0.44	0.45		0.65	0.31	5.94	3.93	0.087
13	Pole44	KF562724	CT	17	162–220 <sup>iii</sup>	0.82	0.90		0.84	0.91	10.60	15.09	0.041
14	Pole45	KF562725	TC	7	122–136 <sup>ii</sup>	0.66	0.66		0.64	0.68	5.87	5.61	–0.006
15	Pole46	KF562726	GAT	4	114–123 <sup>iii</sup>	0.20	0.21		0.16	0.25	3.91	2.00	0.023
Average				9.4		0.5	0.597		0.61	0.53	7.02	6.78	0.039 (.01–.06) <sup>g</sup>

<sup>a</sup> For the allele size range of each marker, the following indicates that a universal, 5' tag (see Methods and Hauswaldt et al. 2003) was added to the primer, adding *X* number of base pairs to the PCR product: <sup>i</sup>NED tag: ACCAACCTAGGAAACACAG; <sup>ii</sup>FAM tag: GGAAACAGCTATGACCAT; <sup>iii</sup>VIC tag: CAGTCGGGCGTCATCA; <sup>iv</sup>PET tag: GGCTAGGAAAGGTTAGTGGC.

<sup>b</sup>  $H_o$  and  $H_e$  represent observed and expected heterozygosities for each locus, respectively (both populations combined,  $N = 95$ ).

<sup>c</sup> Estimated null allele frequency from FreeNA (Chapuis and Estoup 2007) is reported for markers with significant deviation from HWE.

<sup>d</sup> Expected heterozygosity for the Peru ( $N = 47$ ) and Baja ( $N = 48$ ) populations separately.

<sup>e</sup> Allelic richness, estimated with rarefaction in PSTAT (Goudet 1995) for the Peru ( $N = 47$ ) and Baja ( $N = 48$ ) populations.

<sup>f</sup>  $F_{ST}$  was calculated as described by Weir and Cockerham (1984).

<sup>g</sup> Overall  $F_{ST}$  numbers in parentheses represent the lower and upper bounds of the 95% CI obtained via bootstrapping. Markers that deviated from HWE were not included in the population genetic analysis.

di- and hexa-nucleotide repeats (both at 18%), whereas mononucleotide repeats were found at the lowest frequency, just 2.7% (Figure 1B). Looking across species in Figure 1B, it is quite clear that there is substantial variation in the frequencies of repeat classes, even among the related arthropods. Though overall repeat densities were fairly similar between *D. pulex* and *P. elegans*, the distribution of repeat class frequencies were not—patterns for *P. elegans* resembled *D. melanogaster* to a much greater degree than *D. pulex* (Figure 1B). Though broad taxon-specific variation in repeat frequency has been observed (e.g., Tóth et al. 2000), the discrepancy between these crustaceans likely reflects the complexity of repeat evolution (Ellergrén 2004; Meglécz et al. 2012a) and the hundreds of million years of divergence between these species (e.g., Regier et al. 2005).

### Microsatellite Screening, Primer Design, and Primer Testing

Perfect microsatellite sequences were identified for marker development by screening the unique sequences for di-, tri-, tetra-, penta-, and hexa-nucleotide repeats using QDD2.1. In total, we identified 2922 microsatellites of at least 5 repeat units. Primers were successfully designed for 1471 microsatellites (48%), and 45 primer pairs were chosen for testing and further characterization using PCR. In total, 24 of the 45 primers designed produced PCR products with reliable amplification, 15 of which were polymorphic and are

characterized here. A total of 17 primer pairs demonstrated poor amplification, produced PCR products that had more than 2 alleles per individual, or showed unreliable amplification across individuals. Only 4 primer pairs produced no amplification whatsoever. Table 2 displays the results for the 15 successfully designed primer pairs. None of the isolated markers amplified reliably in *P. pollicipes*, *P. polymerus*, or *P. caboverdensis*.

Across the 15 microsatellites, the number of alleles per locus varied from 2 to 22, with an average of 9.4. Observed and expected heterozygosities averaged 0.501, and 0.590, respectively. Most markers consisted of dinucleotide repeats ( $n = 9$ ), but tri-, tetra-, and hexa-nucleotide repeats were also represented in the successfully developed loci (Table 2). One pair of markers, Pole01 and Pole07, showed evidence of significant LD ( $P < 0.00041$ ); the remaining markers showed no significant LD ( $P > 0.07$ ). Four markers exhibited significant deviations from HWE, notably in the direction of fewer heterozygotes than expected (Table 2). Testing these markers for null alleles with FreeNA (Chapuis and Estoup 2007), we found that heterozygote deficiency was likely a result of null alleles, present at low frequency in all 4 of these markers (Table 2). Null alleles are reported often in population genetic literature (reviewed by Dakin and Avise 2004) and are even more common in microsatellite-based analyses of marine invertebrates (e.g., Hare et al. 1996; Hedgecock et al. 2004; Brownlow et al. 2008; Bailie et al. 2010; McInerney

et al. 2011). The low level of null alleles detected here is similar to what has been reported in a population genetic study of the acorn barnacle *Balanus balanoides* (Flight et al. 2012).

### Analysis of Genetic Diversity and Population Structure

With the 11 microsatellite loci that demonstrated HWE, we examined genetic diversity and population structure between the Baja (California, Mexico) and Islilla (Peru) populations. Indices of genetic diversity were consistently lower in the Peruvian population, with 9 of the 11 loci showing lower allelic richness (Sign test,  $P = 0.0654$ ) and 8 of the 11 loci showing lower gene diversity (Wilcoxon signed-rank test,  $P = 0.053$ ; Table 2). The pronounced disparity in diversity could be associated with increased fishing pressure in Peru, but other explanations are likely, including differences in colonization history and effective population size. High variability in the abundance and distribution of *P. elegans* populations in Peru has been reported and is thought to be influenced by strong, episodic El Niño events that disrupt recruitment patterns and affect food availability (Kameya and Zeballos 1988). Substantial fluctuations in population sizes would likely have a large and negative effect on genetic diversity and effective population size in the Peruvian population; however, more complex analyses of effective size and colonization history are required to investigate this result further.

We also found relatively low but significant genetic differentiation between the 2 populations ( $F_{ST} = 0.039$ , 95% CI 0.016–0.061; Table 2). This level of differentiation is a bit lower than expected given the relatively high (1.2%) CO1 divergence and the near reciprocal monophyly of CO1 haplotypes reported by Van Syoc (1994). However,  $F_{ST}$  varied substantially among loci, with some loci showing 0 differentiation (e.g., Pole01 and Pole25) and others exhibiting much higher differentiation (max  $F_{ST} = 0.097$ , Pole29). The relatively low level of overall differentiation suggests that there may be some gene flow between the populations despite the 4000-km distance that separates them. Alternatively, gene flow may be restricted, but not enough time has elapsed for greater divergence in allele frequencies to have accumulated via genetic drift.

### Conclusions

We used a simple and efficient protocol with NGS data to identify and characterize 15 novel polymorphic microsatellite markers for the Pacific gooseneck barnacle *P. elegans*, the first markers of this type developed in the genus *Pollicipes*. These markers had a high rate of amplification success, showed relatively high polymorphism, and revealed differences in genetic diversity and population structure in a preliminary population genetic analysis of 2 populations. Analysis of repeat density demonstrated that *P. elegans* had a fairly high density of microsatellites for its genome size and that there was high variability in repeat class frequency among related taxa, highlighting the complex nature of microsatellite evolution. These markers will be useful for future studies of multiple paternity and

parentage, and a more complete, range-wide investigation of population genetic structure and diversity in *P. elegans*.

### Funding

National Science Foundation grant OCE-0961996 to P.B.M.

### Data Accessibility Policy

The 454 reads are archived in the sequence read archive (accession SRP029161), sequences used to design the 15 microsatellite primers have been assigned GenBank numbers (KF562712–KF562726), and microsatellite data are deposited in the Dryad repository: doi:10.5061/dryad.b7855.

### Acknowledgments

We thank S. Hunicke-Smith of the University of Texas at Austin Genome Sequencing and Analysis Facility for preparation of genomic libraries for 454 sequencing, M. Williams for initial primer screening and testing, S. Bingham and the Arizona State University DNA laboratory for microsatellite sequencing, C. Mayer for kindly providing repeat density data for comparison with *P. elegans*, R. Van Syoc and J. Fernandes for providing specimens of *P. pollicipes*, *P. polymerus*, and *P. caboverdensis* for testing of microsatellite cross amplification, and 2 anonymous reviewers for their helpful comments on previous versions of this manuscript.

### References

- Abdelkrim J, Robertson BC, Stanton JL, Gemmill NJ. 2009. Fast, cost effective development of species-specific microsatellite markers by genomic sequencing. *Biol Tech.* 46:185–191.
- Baillie DA, Fletcher H, Prodöhl PA. 2010. High incidence of cryptic repeated elements in microsatellite flanking regions of galatheid genomes and its practical implications for molecular marker development. *J Crust Biol.* 30:664–672.
- Balzer S, Malde K, Grohme MA, Jonassen I. 2013. Filtering duplicate reads from 454 pyrosequencing data. *Bioinformatics.* 29:830–836.
- Barnes M. 1996. Pedunculate cirripedes of the genus *Pollicipes*. *Oceanogr Mar Biol.* 34:303–394.
- Belkhir K, Borsa P, Chikhi L, Raufaste N, Bonhomme F. 2004. GENETIX 4.05, logiciel sous Windows TM pour la génétique des populations. Laboratoire Génome, Populations, Interactions, CNRS UMR 5171. Montpellier (France): Université de Montpellier II. Available from: <http://www.genetix.univmontp2.fr/genetix/intro.htm>
- Bernard FR. 1988. Potential fishery for the gooseneck barnacle *Pollicipes polymerus* (Sowerby, 1833) in British Columbia. *Fish Res.* 6:287–298.
- Bishop JD, Pemberton AJ. 2006. The third way: spermcast mating in sessile marine invertebrates. *Integr Comp Biol.* 46:398–406.
- Blankenberg D, Gordon A, Von Kuster G, Coraor N, Taylor J, Nekrutenko A; Galaxy Team. 2010. Manipulation of FASTQ data with Galaxy. *Bioinformatics.* 26:1783–1785.
- Brownlow RJ, Dawson DA, Horsburgh GJ, Bell JJ, Fish JD. 2008. A method for genotype validation and primer assessment in heterozygote-deficient species, as demonstrated in the prosobranch mollusc *Hydrobia ulvae*. *BMC Genet.* 9:55.
- Campo D, Molares J, García L, Fernandez-Rueda P, Garcia-Gonzalez C, Garcia-Vazquez E. 2010. Phylogeography of the European stalked barnacle (*Pollicipes pollicipes*): identification of glacial refugia. *Mar Biol.* 157:147–156.
- Chapuis MP, Estoup A. 2007. Microsatellite null alleles and estimation of population differentiation. *Mol Biol Evol.* 24:621–631.

- Dakin EE, Avise JC. 2004. Microsatellite null alleles in parentage analysis. *Heredity*. 93:504–509.
- Doyle JJ, Dickson E. 1987. Preservation of plant samples for DNA restriction endonuclease analysis. *Taxon*. 36:715–722.
- Ellergren H. 2004. Microsatellites: simple sequences with complex evolution. *Nat Rev Genet*. 5:435–445.
- Fernandes JN, Cruz T, Van Syoc R. 2010. *Pollicipes caboverdensis* sp. nov. (Crustacea: Cirripedia: Scalpelliformes), an intertidal barnacle from the Cape Verde Islands. *Zootaxa*. 2557:29–38.
- Flight PA, O'Brien MA, Schmidt PS, Rand DM. 2012. Genetic structure and the North American postglacial expansion of the barnacle, *Semibalanus balanoides*. *J Hered*. 103:153–165.
- Gardner MG, Fitch AJ, Bertozzi T, Lowe AJ. 2011. Rise of the machines—recommendations for ecologists when using next generation sequencing for microsatellite development. *Mol Ecol Resour*. 11:1093–1101.
- Goudet J. 1995. FSTAT (version 1.2): a computer program to calculate F-statistics. *J Hered*. 86:485–486.
- Gregory TR. 2012. Animal genome size database. Available from: <http://www.genomesize.com>
- Hamilton MB, Pincus EL, Di Fiore A, Fleischer RC. 1999. Universal linker and ligation procedures for construction of genomic DNA libraries enriched for microsatellites. *Biotechniques*. 27:500–2, 504.
- Hare MP, Karl SA, Avise JC. 1996. Anonymous nuclear DNA markers in the American oyster and their implications for the heterozygote deficiency phenomenon in marine bivalves. *Mol Biol Evol*. 13:334–345.
- Hauswaldt JS, Glenn TC. 2003. Microsatellite DNA loci from the Diamondback terrapin (*Malaclemys terrapin*). *Mol Ecol Notes*. 3:174–176.
- Hedgecock D, Li G, Hubert S, Bucklin K, Ribes V. 2004. Widespread null alleles and poor cross-species amplification of microsatellite DNA loci cloned from the Pacific oyster, *Crassostrea gigas*. *J Shellfish Res*. 23:379–385.
- Jin L, Baskett ML, Cavalli-Sforza LL, Zhivotovsky LA, Feldman MW, Rosenberg NA. 2000. Microsatellite evolution in modern humans: a comparison of two data sets from the same populations. *Ann Hum Genet*. 64:117–134.
- Kameya A, Zeballos J. 1988. Distribución y densidad de percebes *Pollicipes elegans* (Crustacea: Cirripedia) en el Mediollitoral peruano (Yasla, Paita, Chilca, Lima). *Bol Inst mar Perú-Calloa*. 12:1–22.
- Kelly MW, Grosberg RK, Sanford E. 2012. Love the one you're with: proximity determines paternity success in the barnacle *Tetraclita rubescens*. *Mol Ecol*. 21:5088–5097.
- Liu N, Chen L, Wang S, Oh C, Zhao H. 2005. Comparison of single-nucleotide polymorphisms and microsatellites in inference of population structure. *BMC Genet*. 6(Suppl 1):S26.
- Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, Berka J, Braverman MS, Chen YJ, Chen Z, et al. 2005. Genome sequencing in microfabricated high-density picolitre reactors. *Nature*. 437:376–380.
- Mayer C, Leese F, Tollrian R. 2010. Genome-wide analysis of tandem repeats in *Daphnia pulex*—a comparative approach. *BMC Genomics*. 11:277.
- McInerney CE, Allcock AL, Johnson MP, Bailie DA, Prodöhl PA. 2011. Comparative genomic analysis reveals species-dependent complexities that explain difficulties with microsatellite marker development in molluscs. *Heredity*. 106:78–87.
- Megléc E, Costedoat C, Dubut V, Gilles A, Malausa T, Pech N, Martin JF. 2010. QDD: a user-friendly program to select microsatellite markers and design primers from large sequencing projects. *Bioinformatics*. 26:403–404.
- Megléc E, Nève G, Biffin E, Gardner MG. 2012a. Breakdown of phylogenetic signal: a survey of microsatellite densities in 454 shotgun sequences from 154 non model eukaryote species. *PLoS One*. 7:e40861.
- Megléc E, Pech N, Gilles A, Martin JF, Gardner MG. 2012b. A shot in the genome: how accurately do shotgun 454 sequences represent a genome? *BMC Res Notes*. 5:259.
- Molares J, Freire J. 2003. Development and perspectives for community-based management of the goose barnacle (*Pollicipes pollicipes*) fisheries in Galicia (NW Spain). *Fish Res*. 65:485–492.
- Nei M. 1987. *Molecular evolutionary genetics*. New York: Columbia University Press.
- Niu B, Fu L, Sun S, Li W. 2010. Artificial and natural duplicates in pyrosequencing reads of metagenomic data. *BMC Bioinformatics*. 11:187.
- Ohashi J, Tokunaga K. 2003. Power of genome-wide linkage disequilibrium testing by using microsatellite markers. *J Hum Genet*. 48:487–491.
- Quinteiro J, Rodríguez-Castro J, Lopez P, Lopez-Jurado LF, Gonzalez-Henriquez N, Rey-Mendez M. 2011. Morphological and molecular characterization of a new species of Atlantic stalked barnacle (Scalpelliformes: Pollicipedidae) from the Cape Verde Islands. *J Mar Biol Assoc UK*. 91:1537–1545.
- Quinteiro J, Rodríguez-Castro J, Rey-Méndez M. 2007. Population genetic structure of the stalked barnacle *Pollicipes pollicipes* (Gmelin, 1789) in the northeastern Atlantic: influence of coastal currents and mesoscale hydrographic structures. *Mar Biol*. 153:47–60.
- Regier JC, Shultz JW, Kambic RE. 2005. Pancrustacean phylogeny: hexapods are terrestrial crustaceans and maxillopods are not monophyletic. *Proc Biol Sci*. 272:395–401.
- Rosenberg NA, Li LM, Ward R, Pritchard JK. 2003. Informativeness of genetic markers for inference of ancestry. *Am J Hum Genet*. 73:1402–1422.
- Tóth G, Gáspári Z, Jurka J. 2000. Microsatellites in different eukaryotic genomes: survey and analysis. *Genome Res*. 10:967–981.
- Van Syoc RJ. 1994. Genetic divergence between subpopulations of the eastern Pacific goose barnacle *Pollicipes elegans*: mitochondrial cytochrome c subunit 1 nucleotide sequences. *Mol Mar Biol Biotechnol*. 3:338–346.
- Veliz D, Duchesne P, Bourget E, Bernatchez L. 2006. Genetic evidence for kin aggregation in the intertidal acorn barnacle (*Semibalanus balanoides*). *Mol Ecol*. 15:4193–4202.
- Wang J, Santure AW. 2009. Parentage and sibship inference from multilocus genotype data under polygamy. *Genetics*. 181:1579–1594.
- Weir BS, Cockerham CC. 1984. Estimating F-statistics for the analysis of population structure. *Evolution*. 38:1358–1370.
- Zane L, Bargelloni L, Patarnello T. 2002. Strategies for microsatellite isolation: a review. *Mol Ecol*. 11:1–16.

Received January 15, 2013; First decision March 18, 2013;  
Accepted August 28, 2013

Corresponding Editor: Jose Lopez